

Minimal Matrix Representations of Five-Dimensional Lie Algebras

R. GHANAM, G. THOMPSON

*Department of Mathematics, Virginia Commonwealth University in Qatar,
PO Box 8095, Doha, Qatar
raghanam@vcu.edu*

*Department of Mathematics, University of Toledo,
Toledo, OH 43606, U.S.A.
gerard.thompson@utoledo.edu*

Presented by Juan Carlos Marrero

Received July 9, 2014

Abstract: We obtain minimal dimension matrix representations for each indecomposable five-dimensional Lie algebra over \mathbb{R} and justify in each case that they are minimal. In each case a matrix Lie group is given whose matrix Lie algebra provides the required representation.

Key words: Lie algebra, Lie group, minimal representation.

AMS *Subject Class.* (2010): 17B, 53C.

1. INTRODUCTION

Given a real Lie algebra \mathfrak{g} of dimension n a well known theorem due to Ado asserts that \mathfrak{g} has a faithful representation as a subalgebra of $gl(p, \mathbb{R})$ for some p . The theorem does not give much information about the value of p but leads one to believe that p may be very large in relation to the size of n and consequently it seems to be of limited practical value. We define the invariant $\mu(\mathfrak{g})$ to be the minimum value of p . A little care must be exercised because there may well be inequivalent representations for which this minimum value is attained. Of course if \mathfrak{g} has a trivial center then the adjoint representation furnishes a faithful representation of \mathfrak{g} and in the notation used above $\mu(\mathfrak{g}) \leq n$. Nonetheless many algebras have non-trivial centers, nilpotent algebras for example, and then the adjoint representation is not faithful. Even if the center is trivial it could well be the case that $\mu(\mathfrak{g}) < n$.

Of course it is interesting to ascertain the value of μ from a theoretical point of view. However, an important practical reason is that calculations involving symbolic programs such as Maple and Mathematica use up lots of

memory when storing matrices; accordingly, calculations are likely to be faster if one can represent matrix Lie algebras using matrices of a small size.

In two recent papers [1, 5], the problem of finding such a minimal representation is considered for the four-dimensional Lie algebras. In [5], the main technique is somewhat indirect and depends on a construction known as a left symmetric structure. In [1], the minimal representation has been calculated directly without the need for considering left symmetric structures.

In this paper, we consider all indecomposable five-dimensional Lie algebras listed in [7]. It forms part of a series in which we hope to find minimal dimensional representations for all the low-dimensional algebras. Partial classifications of Lie algebras are known up to dimension nine. The five-dimensional problem is of a completely magnitude from the four-dimensional case since there are, up to isomorphism, forty classes of algebra. It has already been shown in [2] that $\mu(\mathfrak{g}) \leq 5$ for each five-dimensional Lie algebra in the list. In this paper we obtain sharper results and for the cases where $\mu(\mathfrak{g}) < 5$ we give an explicit representation for the Lie algebra. This representation is given by means of a matrix Lie group, denoted by S , in local coordinates (w, x, y, z, q) . We also explain for those cases where $\mu(\mathfrak{g}) = 5$ why we cannot have $\mu(\mathfrak{g}) < 5$. The matrix S can be modified by adding quadratic and higher order terms to the entries without affecting the matrix Lie algebra, obtained by differentiating and evaluating at the identity. Accordingly, as a check, we supply the corresponding right-invariant vector fields. We also adopt the convention that when referring to an “abstract” Lie algebra five-dimensional we use $\{e_1, e_2, e_3, e_4, e_5\}$ as a basis but when we construct a matrix representation we use $\{E_1, E_2, E_3, E_4, E_5\}$ for the corresponding generators.

An outline of the paper is as follows. In Section 2 we give a brief overview of the indecomposable five-dimensional Lie algebras. In Section 3 we give two results about representations that are needed in the sequel. We shall also have occasion to use a version of Lie’s Theorem for Lie algebras over \mathbb{R} . In Section 4 we determine the few algebras where $\mu = 4$ as result of the representing matrices having some partial complex structure. In Section 5 we are able to determine all algebras for which $\mu = 3$. In Section 6 we consider algebras that have a four-dimensional abelian nilradical and in Section 7 the six-dimensional nilpotent algebras. In Section 8 we give a group matrix S for each of the five-dimensional Lie algebras. In fact for the convenience of the reader we provide also an S -matrix for all indecomposable Lie algebras of dimension *and less*. We are able to reduce the problem to the consideration of five difficult cases, namely, $A_{5,21}$, $A_{5,22}$, $A_{5,23(b \neq 1)}$, $A_{5,31}$, $A_{5,38}$. These five cases are relegated to

an appendix which consists of a lot of technical details. For the casual reader Section 8 is likely to be of most interest. Nonetheless, for the sake of the integrity of the results, we feel that it is essential to supply these details so that they can be verified independently. Following [7] we denote each of the five-dimensional algebras as $A_{5,k}$ where $1 \leq k \leq 40$.

We close this Introduction by stating several conclusions. First of all one might imagine that many algebras could have representations with $\mu = 4$ where the representing matrices having some partial complex structure as necessitated by applying Lie's Theorem for Lie algebras over \mathbb{R} rather than over \mathbb{C} . However, relatively few such representations actually occur. Secondly, and despite the first remark, for many of the five-dimensional indecomposable Lie algebras we do have $\mu = 4$. Some of the algebras for which $\mu = 5$ are *CR*-Lie algebras and over \mathbb{C} we would have $\mu = 4$, for example $A_{5,25}$ and $A_{5,26}$ which are equivalent over \mathbb{C} to $A_{5,19}$ and $A_{5,20}$, respectively. Also, there are some classes of algebra depending on parameters where $\mu(\mathfrak{g}) < 5$ *but only for certain values of the parameters*. In fact, as a crude count, 21 of the 40 algebras have $\mu = 5$, although $\mu(\mathfrak{g}) < 5$ for several of those 21 algebras for special values of the parameters. In a separate paper we shall construct minimal dimension matrix representations for *decomposable* five-dimensional Lie algebras. Finally, in this paper the many calculations were performed with the help of the symbolic manipulation program Maple.

2. REAL AND COMPLEX FIVE-DIMENSIONAL LIE ALGEBRAS

The *real* five-dimensional indecomposable Lie algebras were classified by G. Mubarakzyanov [6]. They can be found easily in [7] and we list them in Section 4. The first six algebras are nilpotent. These six algebras are distinguished by their index of nilpotence and the dimension of the derived algebra except for the filiforms $A_{5,2}$ and $A_{5,6}$. However, $A_{5,2}$ has a codimension one abelian ideal, whereas $A_{5,6}$ does not so the six algebras are mutually non-isomorphic.

We remark that the algebras $A_{5,7} - A_{5,18}$ have a four-dimensional abelian nilradical; $A_{5,19} - A_{5,29}$ have a four-dimensional non-abelian nilradical isomorphic to $H \oplus \mathbb{R}$, where H denotes the three-dimensional Heisenberg algebra; $A_{5,30} - A_{5,32}$ have a nilradical that is isomorphic to the unique four-dimensional indecomposable nilpotent algebra; $A_{5,33} - A_{5,35}$, $A_{5,38}$, $A_{5,39}$ have an abelian three-dimensional nilradical and $A_{5,36}$, $A_{5,37}$ have a three-dimensional nilradical that is isomorphic to H ; $A_{5,40}$ is the only algebra that is not solvable,

that is, has a non-trivial Levi decomposition, being a semi-direct product of $\mathfrak{sl}(2, \mathbb{R})$ and the abelian algebra \mathbb{R}^2 . In particular, all of these algebras, with the exception of $A_{5,36}$, $A_{5,37}$ and $A_{5,40}$ have a three-dimensional abelian subalgebra.

Given a Lie algebra \mathfrak{g} we denote its derived algebra by $[\mathfrak{g}, \mathfrak{g}]$. If now \mathfrak{g} is an indecomposable five-dimensional Lie algebra the dimension of $[\mathfrak{g}, \mathfrak{g}]$ is precisely one only for the Heisenberg algebra $A_{5,4}$ and is two in just $A_{5,1}$ and $A_{5,5}$ which are both nilpotent. *Apart from the non-solvable $A_{5,40}$, for all other indecomposable five-dimensional indecomposable algebras the dimension of $[\mathfrak{g}, \mathfrak{g}]$ is either three or four.* In fact in precisely the following algebras the dimension of $[\mathfrak{g}, \mathfrak{g}]$ is three: $A_{5,2}$, $A_{5,3}$, $A_{5,6}$, $A_{5,8}$, $A_{5,10}$, $A_{5,14}$, $A_{5,15(a=0)}$, $A_{5,19(a=1)}$, $A_{5,20(a=0,1)}$, $A_{5,22}$, $A_{5,26(p=0)}$, $A_{5,27}$, $A_{5,28(a=1)}$, $A_{5,29}$, $A_{5,30(a=1)}$, $A_{5,32} - A_{5,39}$.

3. TWO REPRESENTATION RESULTS

3.1. Transposition about the anti-diagonal.

PROPOSITION 3.1. *Suppose that the Lie algebra \mathfrak{g} has a representation as a subalgebra of $\mathfrak{gl}(p, \mathbb{R})$. Suppose that $L : \mathfrak{gl}(p, \mathbb{R}) \rightarrow \mathfrak{gl}(p, \mathbb{R})$ is a (linear) involution, that is, has period two. Then mapping a representing matrix M to $-LM^tL$ gives a second inequivalent representation of \mathfrak{g} .*

Proof. Given $M \in \mathfrak{gl}(p, \mathbb{R})$ a representing matrix for \mathfrak{g} , map it to $\phi(M) = -LM^tL$. For a second such matrix N we have

$$\begin{aligned} [\phi(M), \phi(N)] &= [-LM^tL, -LN^tL] = [LM^tLLN^tL - LN^tLLM^tL] \\ &= [LM^tN^tL - LN^tM^tL] = L[N, M]^tL = \phi([M, N]). \end{aligned}$$

■

COROLLARY 3.2. *Suppose that the Lie algebra \mathfrak{g} has a representation as a subalgebra of $\mathfrak{gl}(p, \mathbb{R})$. Then transposing the representing matrices about the anti-diagonal (and taking negatives) gives a second inequivalent (that is not necessarily equivalent) representation of \mathfrak{g} .*

Proof. Take for L in the Proposition the matrix whose only non-zero entries are 1's down the anti-diagonal. Then the map ϕ consists of taking a negative and transposing about the anti-diagonal. ■

many cases for representing five-dimensional algebras in $\mathfrak{gl}(4, \mathbb{R})$. However, it is remarkable that cases (i-iv) correspond to representations for only a very limited number of and, in some cases, unexpected algebras. In the interests of concision we shall be content merely to sketch the results.

Notice first of all that it follows from Section 3.3 that if a five-dimensional algebra \mathfrak{g} has a representation of type (i)-(iv) then at least one of its adjoint matrices must have non-real eigenvalues. Actually this remark is not obvious but is nonetheless true as we shall explain. In any case given the last remark, we can proceed as follows. First of all, the nilpotent algebras $A_{5,1} - A_{5,6}$ cannot have representations of types (i)-(iv). Secondly, in Section 6 we give a systematic analysis of algebras $A_{5,7} - A_{5,18}$ so we leave them to one side for the moment. Thirdly, of algebras $A_{5,19} - A_{5,39}$ only $A_{5,25}$, $A_{5,26}$, $A_{5,35}$, $A_{5,37}$ and $A_{5,39}$ have adjoint matrices with non-real eigenvalues. As regards $A_{5,35}$ we know that $\mu = 4$ in view of Theorem 3.1. Similarly we will show that $\mu = 4$ for $A_{5,37}$ in Section 5. As for $A_{5,39}$ it is equivalent as a complex algebra to $A_{5,38}$ and the latter will be shown to have $\mu = 5$ so that $\mu = 5$ for $A_{5,39}$ also. Hence of algebras $A_{5,19} - A_{5,39}$ whose adjoint matrices have non-real eigenvalues, μ is in doubt only for $A_{5,25}$ and $A_{5,26}$.

PROPOSITION 4.1. *For $A_{5,25}$ and $A_{5,26}$ we have $\mu = 5$.*

Proof. Assuming that $\mu = 4$ we shall obtain a contradiction for each of cases (i-v). First of all case (v) is excluded because all of the adjoint matrices would have real eigenvalues which is not true for $A_{5,25}$ and $A_{5,26}$. As regards case (i) for a Lie algebra \mathfrak{g} , the derived algebra $[\mathfrak{g}, \mathfrak{g}]$ must consist of matrices of the form

$$\begin{bmatrix} 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

where only the asterisks can be non-zero. It must be abelian and hence \mathfrak{g} cannot be $A_{5,25}$ or $A_{5,26}$. Case (iii) can be excluded in view of Proposition 3.1 once we have ruled out case (ii) which we look at next. The derived algebra must consist of matrices of the form

$$\begin{bmatrix} 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

where only the asterisks can be non-zero. Let us suppose first of all that we have one non-nilpotent matrix E of type (ii). Then we obtain a six-dimensional solvable algebra that has the nilpotent algebra $A_{5,1}$ as its nilradical and E spans a complement to it. In this six-dimensional algebra we find that $\text{ad } E$ has eigenvalues $\{0, g - i, a - i \pm b\sqrt{-1}, a - g \pm b\sqrt{-1}\}$. The restriction of $\text{ad } E$ to the five-dimensional subalgebra must have eigenvalues that are a subset of these six and of course 0 must be one of them. The only possibility is to remove $g - i$ which leaves the eigenvalues $\{0, a - i \pm b\sqrt{-1}, a - g \pm b\sqrt{-1}\}$ and we can only have $A_{5,17prs}$ or $A_{5,18}$. In fact $A_{5,17prs(s=1)}$ and $A_{5,18}$ do occur and examples of each are listed in Section 10. However, $A_{5,25}$ and $A_{5,26}$ are excluded because their ad -matrices have at most a pair of complex eigenvalues and not two pairs.

If there are two non-nilpotent matrices of type (ii), E_6 and E_7 say, then $[\mathfrak{g}, \mathfrak{g}]$ is three-dimensional which as regards $A_{5,25}$ and $A_{5,26}$ would only allow $A_{5,26}, p = 0$. We shall obtain together with the representation of $A_{5,1}$ a seven-dimensional codimension two nilradical solvable algebra and the question now is whether there exists a five-dimensional subalgebra besides $A_{5,1}$. Such an algebra \mathfrak{g} must have $[\mathfrak{g}, \mathfrak{g}]$ of dimension three. Again letting E_6 be the given matrix of type (ii) in 1 we find that the eigenvalues of $\text{ad } E$ are $\{0, 0, i - g, a - g \pm b\sqrt{-1}, a - i \pm b\sqrt{-1}\}$. Now comparing with the adjoint matrices of $A_{5,26}, p = 0$ we deduce that $a = g = i, b = 1$. In fact arbitrary multiples of the identity can be added to E_6 and E_7 without affecting the putative representation that we are looking for. However the same argument that applies to E_6 applies equally to E_7 which would imply that a linear combination of E_6 and E_7 is nilpotent, a contradiction.

Case (iv) is similar to case (ii). Now $[\mathfrak{g}, \mathfrak{g}]$ consist of matrices of the form

$$\begin{bmatrix} 0 & * & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

where only the asterisks can be non-zero. Let us suppose first of all that we have one non-nilpotent matrix E of type (iv). Then we obtain a six-dimensional solvable algebra that has the nilpotent algebra $A_{5,4}$ as its nilradical and E spans a complement to it. However, in the latter algebra we find that $\text{ad } E$ has eigenvalues $\{0, a - i, a - e \pm f\sqrt{-1}, e - i \pm f\sqrt{-1}\}$. The restriction of $\text{ad } E$ to \mathfrak{g} must have eigenvalues that are a subset of these six and again we find two pairs of non-real complex conjugates.

If there are two non-nilpotent matrices E_6 and E_7 of type (iv) we shall obtain together with the representation of $A_{5,4}$ a seven-dimensional codimension two nilradical solvable algebra. Such an algebra \mathfrak{g} must have $[\mathfrak{g}, \mathfrak{g}]$ of dimension three. Once again we find that, matching adjoint matrices to $A_{5,26}, p = 0$, a linear combination of E_6 and E_7 is nilpotent, a contradiction. ■

5. REPRESENTATIONS IN $\mathfrak{gl}(3, \mathbb{R})$

Let \mathfrak{g} be an indecomposable real solvable five-dimensional algebra. Clearly \mathfrak{g} cannot have a representation as a subalgebra of $\mathfrak{gl}(2, \mathbb{R})$ so the smallest value of n for which \mathfrak{g} can be represented in $\mathfrak{gl}(n, \mathbb{R})$ is $n = 3$.

LEMMA 5.1. *Any three-dimensional abelian subalgebra of $\mathfrak{gl}(3, \mathbb{R})$ contains a multiple of the identity.*

Proof. Consider an element of such an abelian subalgebra. We can put it into one of the following four forms by a real change of basis:

$$(a) \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{bmatrix}, \quad (b) \begin{bmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 1 \\ 0 & 0 & \lambda \end{bmatrix}, \quad (c) \begin{bmatrix} \lambda & 0 & 1 \\ 0 & \mu & 0 \\ 0 & 0 & \lambda \end{bmatrix}, \quad (d) \begin{bmatrix} \alpha & \beta & 0 \\ -\beta & \alpha & 0 \\ 0 & 0 & \gamma \end{bmatrix} \quad (\beta \neq 0).$$

The proof follows easily now by considering the centralizer in each of these four cases. ■

COROLLARY 5.2. *The only solvable five-dimensional indecomposable algebras that could be represented as a subalgebra of $\mathfrak{gl}(3, \mathbb{R})$ are $A_{5,36}$ or $A_{5,37}$.*

Proof. If a subalgebra of $\mathfrak{gl}(3, \mathbb{R})$ contains I it will be decomposable. However, the only algebras that have do not have an abelian three-dimensional subalgebra are $A_{5,36}$ and $A_{5,37}$. ■

As regards $A_{5,37}$ let us note that it is equivalent over \mathbb{C} to $A_{5,36}$ which means that both are equivalent considered as algebras over \mathbb{C} . To see how, make a change of basis according to

$$\begin{aligned} \bar{e}_1 &= -\frac{e_1}{2}, & \bar{e}_2 &= \frac{e_2 + e_3}{2}, & \bar{e}_3 &= \frac{\sqrt{-1}(e_2 - e_3)}{2}, \\ \bar{e}_4 &= 2e_4 + e_5, & \bar{e}_5 &= -\sqrt{-1}e_5. \end{aligned}$$

We obtain the following brackets, which are formally identical to $A_{5,36}$ except for the factor of $\sqrt{-1}$ which can then be removed by scaling e_1 :

$$\begin{aligned} [e_2, e_3] &= \sqrt{-1}e_1, & [e_1, e_4] &= e_1, & [e_2, e_4] &= e_2, \\ [e_2, e_5] &= -e_2, & [e_3, e_5] &= e_3. \end{aligned}$$

The Lie algebra $A_{5,36}$ does have a representation in $\mathfrak{gl}(3, \mathbb{R})$: it is isomorphic to the space of trace-free upper triangular matrices.

Since $A_{5,37}$ is equivalent over \mathbb{C} to $A_{5,36}$ it certainly has a representation in $\mathfrak{gl}(3, \mathbb{C})$. However, we claim that $A_{5,37}$ does not have a representation in $\mathfrak{gl}(3, \mathbb{R})$. Indeed if it did by Lie's Theorem it would have a representation by matrices of the form

$$\begin{bmatrix} a & b & c \\ -b & a & d \\ 0 & 0 & e \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} c & d & e \\ 0 & a & b \\ 0 & -b & a \end{bmatrix}.$$

However, in either case the derived algebra is at most two-dimensional whereas for $A_{5,37}$ it is three-dimensional so $A_{5,37}$ cannot be represented in $\mathfrak{gl}(3, \mathbb{R})$.

We remark finally that algebra $A_{5,40}$ is the Lie algebra of the special affine group and so by its very definition has a representation in $\mathfrak{gl}(3, \mathbb{R})$ but not an upper triangular representation. Thus we have determined those algebras that can be represented in $\mathfrak{gl}(3, \mathbb{R})$ and $\mathfrak{gl}(3, \mathbb{C})$.

6. FOUR-DIMENSIONAL ABELIAN NILRADICAL ALGEBRAS

Now we consider algebras $\mathfrak{g} A_{5,7} - A_{5,18}$ for which the algebra is solvable but not nilpotent and for which $\text{nil}(\mathfrak{g})$ is abelian. We quote next a result of Schur-Jacobson [4].

PROPOSITION 6.1. (SCHUR-JACOBSON) *The maximal commutative subalgebra of $\mathfrak{gl}(n, \mathbb{R})$ is of dimension $1 + \lfloor \frac{n^2}{4} \rfloor$ where $\lfloor \cdot \rfloor$ denotes the integer part of a real number. Up to change of basis if n is even the subalgebra consists of the upper left hand block with row entries running from $1 + \frac{n}{2}$ to n and column entries from 1 to $\frac{n}{2}$ together with multiples of the identity; if n is odd the subalgebra consists of the upper left hand blocks with row entries running from either $\frac{n+1}{2}$ to n and column entries from 1 to $\frac{n-1}{2}$ or $\frac{n+3}{2}$ to n and 1 to $\frac{n+1}{2}$, respectively, together with multiples of the identity.*

We apply the Proposition in the case $n = 4$. We shall not want to include multiples of the identity because it will lead to a decomposable algebra. Accordingly we assume that we have a basis for $\text{nil}(\mathfrak{g})$ of the following form:

$$E_1 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad E_2 = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$E_3 = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad E_4 = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

To obtain a full basis for \mathfrak{g} we add a generator E_5 of the form

$$E_5 = \begin{bmatrix} a & b & 0 & 0 \\ c & d & 0 & 0 \\ 0 & 0 & e & f \\ 0 & 0 & g & h \end{bmatrix}.$$

This form of E_5 is dictated by the requirement that $\text{ad } E_5$ map $\text{nil}(\mathfrak{g})$ spanned by E_1, E_2, E_3, E_4 to itself. Then the Jacobi identity is satisfied and we have the following brackets:

$$[E_1, E_5] = (h - a)E_1 + gE_2 - cE_3, \quad [E_2, E_5] = fE_1 + (e - a)E_2 - cE_4,$$

$$[E_3, E_5] = -bE_1 + (h - d)E_3 + gE_4, \quad [E_4, E_5] = -bE_2 + fE_3 + (e - d)E_4.$$

Now consider $\text{ad } E_5$. Its bottom row and last column are zero. We look at the upper right 4×4 block given by

$$M = \begin{bmatrix} h - a & f & -b & 0 \\ g & e - a & 0 & -b \\ -c & 0 & h - d & f \\ 0 & -c & g & e - d \end{bmatrix},$$

and consider its possible Jordan normal form: for Lie algebras with abelian codimension one nilradical the Jordan normal form is a complete invariant apart from an overall scaling. The matrix M enjoys a particular property.

LEMMA 6.2. *If $\lambda_1, \lambda_2, \lambda_3, \lambda_4$ are the eigenvalues of M there is an ordering of them so that $\lambda_1 + \lambda_2 = \lambda_3 + \lambda_4$.*

Proof. In fact solving explicitly we find that the eigenvalues are of the form $A \pm \sqrt{B \pm \sqrt{C}}$ where A, B, C are determined in terms of a, b, c, d, e, f, g, h . ■

We compare with algebras $A_{5,7} - A_{5,18}$ and obtain various of these algebras sometimes for special values of the parameters. It is too much to be able to find the Jordan normal forms for the matrix in the form above. However, going back to the the original form of the algebra we can make a change of basis of the form

$$S = \begin{bmatrix} P & 0 \\ 0 & Q \end{bmatrix},$$

where P and Q are arbitrary non-singular matrices. More precisely we multiply each of the matrices E_i ($1 \leq i \leq 5$) on the left by S^{-1} and on the right by S ; although each of the E_i ($1 \leq i \leq 4$) are changed we can in effect use the original E_i as part of the basis since the E_i 's are transformed into linear combinations of themselves. As regards the matrix E_5 , its two blocks are conjugated separately by P and Q . As such we may assume that each of these blocks are in Jordan normal form. There are three Jordan normal forms for a 2×2 matrix giving nine cases in toto. We summarize these cases as follows in which we give first of all the conditions on a, b, c, d, e, f, g, h to be in one of the nine cases, then the eigenvalues λ of the matrix $\text{ad}(E_5)$ and finally conditions on the parameters of the algebras $A_{5,7} - A_{5,18}$ so that they can have a representation in $\mathfrak{gl}(4, \mathbb{R})$:

- (1) $c = -b, d = a, g = -f, h = e$ ($bf \neq 0$): $\lambda = e - a \pm ib \pm if$,
 $A_{5,13apr}$ $g = a = p = 1, f = b = \frac{r}{2}$,
 $A_{5,17}$ $e - a = p = r, b + f = 1, b - f = s$;
- (2) $c = -b, d = a, f = 0, g = 0$ ($b \neq 0$): $\lambda = e - a \pm ib, h - a \pm ib$,
 $A_{5,17prs}$ $e - a = p, h - a = q, b = s = 1$;
- (3) $c = -b, d = a, h = e, f = 1, g = 0$ ($b \neq 0$): $\lambda = e - a \pm ib, e - a \pm ib$,
 $A_{5,18p}$ $e - a = p, b = 1$;
- (4) $c = 0, b = 0, g = -f, h = e$ ($f \neq 0$): $\lambda = e - a \pm if, e - d \pm if$,
 $A_{5,17prs}$ $e - a = p, e - d = q, f = s = 1$;

- (5) $c = 0, b = 0, f = 0, g = 0$: $\lambda = e - a, e - d, h - a, h - d$,
 $A_{5,7abc}$ $e - a = 1, e - d = \bar{a}, h - a = \bar{b}, h - d = \bar{a}$ so $\bar{a} + \bar{b} - \bar{c} = 1$;
- (6) $c = 0, b = 0, h = e, f = 1, g = 0$: $\lambda = e - a, e - a, e - d, e - d$,
 $A_{5,15a}$ $e - a = 1, e - d = \bar{a}$;
- (7) $d = a, c = 0, b = 1, g = -f, h = e$ ($f \neq 0$): $\lambda = e - a \pm if, e - a \pm if$,
 $A_{5,18p}$ $e - a = p, f = 1$;
- (8) $d = a, c = 0, b = 1, f = 0, g = 0$: $\lambda = e - a, e - a, h - a, h - a$,
 $A_{5,15a}$ $e - a = 1, h - a = \bar{a}$;
- (9) $d = a, c = 0, b = 1, h = e, f = 1, g = 0$: $\lambda = e - a, e - a, e - a, e - a$,
 $A_{5,11c}$ $e - a = 1$ so $\bar{c} = 1$.

To summarize: in $A_{5,7} - A_{5,18}$, the algebras $A_{5,8c}, A_{5,9bc}, A_{5,10}, A_{5,12}, A_{5,14p}$ and $A_{5,16pq}$ do not occur at all as subalgebras of $\mathfrak{gl}(4, \mathbb{R})$. Algebras $A_{5,15a}$ and $A_{5,18p}$ do occur and $A_{5,7abc}$ occurs but only for the case $a + b - c = 1$, $A_{5,11c}$ for $c = 1$, $A_{5,13apr}$ for $a = p = 1$ and $A_{5,17}$ for $p = r$ or $s = 1$.

7. FIVE-DIMENSIONAL NILPOTENT ALGEBRAS

Now suppose that \mathfrak{g} is nilpotent and can be represented in $\mathfrak{gl}(4, \mathbb{C})$. We note that both the algebras $A_{5,1}$ and $A_{5,2}$ have a four-dimensional abelian subalgebra indeed ideal. For these algebras we can proceed much as we did for the abelian nilradical case except now we must have that $\text{ad } E_5$ is nilpotent. We note first of all that we can add a multiple of the identity to E_5 and it does not change $\text{ad } E_5$ so we may assume that E_5 has trace zero and so we put $h = -(a + d + e)$. Next we shall demand that $\text{ad } E_5$ has trace zero which gives that $d = -a$ and then we find that the trace of $(\text{ad } E_5)^3$ is zero. In order to make $\text{ad } E_5$ nilpotent it is sufficient to have that the trace of $(\text{ad } E_5)^2$ and the determinant of $\text{ad } E_5$ zero for then $\text{ad } E_5$ will have all eigenvalues zero. These conditions give us $a^2 + bc = 0$ and $e^2 + fg = 0$. In order for E_5 not to vanish entirely and using the block change of basis we can reduce to the cases where only $b = 1$ or $f = 1$ are the only non-zero entries or else $b = f = 1$. The first two of these cases correspond to $A_{5,1}$ whereas the third after making a change of basis

$$\begin{aligned} e'_1 &= \frac{e_1 - e_2}{2}, & e'_2 &= e_2, & e'_3 &= -\frac{e_3}{2}, \\ e'_4 &= e_1 + e_4, & e'_5 &= e_5 \end{aligned}$$

becomes $[e_1, e_5] = e_2$, $[e_3, e_5] = e_1$. This algebra is decomposable being a direct sum of $A_{4,1}$ and \mathbb{R} . In particular for $A_{5,2}$ we cannot have $\mu = 4$.

Now we consider algebras $A_{5,3}$ and $A_{5,6}$. For $A_{5,3}$ we have non-zero brackets $[e_3, e_4] = e_2$, $[e_3, e_5] = e_1$, $[e_4, e_5] = e_3$. The first two brackets give us a copy of $A_{5,1}$; indeed if we permute e_3 and e_5 and then change the sign of e_5 we obtain $[e_3, e_5] = e_1$, $[e_4, e_5] = e_2$, $[e_3, e_4] = e_5$. Now the argument above gave us, up to change of basis, a unique representation of $A_{5,1}$ in $\mathfrak{gl}(4, \mathbb{R})$. Since $[e_3, e_4] \neq e_5$ we conclude that there is no representation of $A_{5,3}$ in $\mathfrak{gl}(4, \mathbb{R})$. Similarly for $A_{5,6}$ $[e_3, e_4] = e_1$, $[e_2, e_5] = e_1$, $[e_3, e_5] = e_2$, $[e_4, e_5] = e_3$ the latter three brackets produce a copy of $A_{5,2}$. However, we have shown that there is no representation of $A_{5,2}$ in $\mathfrak{gl}(4, \mathbb{R})$ and therefore not for $A_{5,6}$, either.

8. GROUP REPRESENTATIONS CORRESPONDING TO LIE ALGEBRAS IN DIMENSION ≤ 5

- $A_{2,1}$ $[e_1, e_2] = e_2$:

$$S = \begin{bmatrix} e^x & y \\ 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $-(D_x + yD_y)$, D_y .

- $A_{3,1}$ $[e_2, e_3] = e_1$:

$$S = \begin{bmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: D_z , D_y , $D_x + yD_z$.

- $A_{3,2}$ $[e_1, e_3] = e_1$, $[e_2, e_3] = e_1 + e_2$:

$$S = \begin{bmatrix} e^z & ze^z & x \\ 0 & e^z & y \\ 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: D_x , D_y , $D_z + (x + y)D_x + yD_y$.

- $A_{3.3}$ ($a = 1$), $A_{3.4}$ ($a = -1$), $A_{3.5a}$ ($0 < |a| < 1$) $[e_1, e_3] = e_1$, $[e_2, e_3] = ae_2$:

$$S = \begin{bmatrix} e^z & 0 & x \\ 0 & e^{az} & y \\ 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, D_z + xD_x + ayD_y$.

- $A_{3.6}$ ($a = 0$), $A_{3.7a}$ ($a > 0$) $[e_1, e_3] = ae_1 - e_2$, $[e_2, e_3] = e_1 + ae_2$:

$$S = \begin{bmatrix} e^{az} \cos z & e^{az} \sin z & x \\ -e^{az} \sin z & e^{az} \cos z & y \\ 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, D_z + (ax + y)D_x + (ay - x)D_y$.

- $A_{3.8}$ $[e_1, e_3] = -2e_2$, $[e_1, e_2] = e_1$, $[e_2, e_3] = e_3$:

$$S = \begin{bmatrix} \cosh x + \sinh x \cosh y & -e^{-z} \sinh x \sinh y \\ e^z \sinh x \sinh y & \cosh x - \sinh x \cosh y \end{bmatrix}$$

Right-invariant vector fields:

$$\begin{aligned} & \frac{e^z}{2} \left(\sinh y D_x - \frac{(\cosh x \cosh y - \sinh x)}{\sinh x} D_y - \frac{(\sinh x \cosh y - \cosh x)}{(\sinh y \sinh x)} D_z \right), \\ & \frac{1}{2} \left(\cosh y D_x - \frac{\cosh x \sinh y}{\sinh x} D_y - D_z \right), \\ & \frac{e^{-z}}{2} \left(-\sinh y D_x + \frac{(\cosh x \cosh y + \sinh x)}{\sinh x} D_y + \frac{(\sinh x \cosh y + \cosh x)}{(\sinh y \sinh x)} D_z \right). \end{aligned}$$

- $A_{3.9}$ $[e_1, e_2] = e_3$, $[e_2, e_3] = e_1$, $[e_3, e_1] = e_2$:

$$S = \begin{bmatrix} \cos x \cos y \cos z - \sin x \sin z & \sin x \cos y \cos z + \cos x \sin z & -\sin y \cos z \\ -\cos x \cos y \sin z - \sin x \cos z & -\sin x \sin z \cos y + \cos x \cos z & \sin y \sin z \\ \cos x \sin y & \sin x \sin y & \cos y \end{bmatrix}.$$

Right-invariant vector fields: $D_z, \frac{\sin z}{\sin y} D_x + \cos z D_y - \frac{\cos y \sin z}{\sin y} D_z, \frac{\cos z}{\sin y} D_x - \sin z D_y - \frac{\cos y \cos z}{\sin y} D_z$.

- $A_{4.1}$ $[e_2, e_4] = e_1$, $[e_3, e_4] = e_2$:

$$S = \begin{bmatrix} 1 & w & \frac{w^2}{2} & x \\ 0 & 1 & w & y \\ 0 & 0 & 1 & z \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, D_z, D_w + yD_x + zD_y$.

- $A_{4.2a}$ ($a \neq 0$) $[e_1, e_4] = ae_1$, $[e_2, e_4] = e_2$, $[e_3, e_4] = e_2 + e_3$:

$$S = \begin{bmatrix} e^{aw} & 0 & 0 & x \\ 0 & e^w & we^w & y \\ 0 & 0 & e^w & z \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, D_z, D_w + axD_x + (y + z)D_y + zD_z$.

- $A_{4.3}$ $[e_1, e_4] = e_1$, $[e_3, e_4] = e_2$:

$$S = \begin{bmatrix} e^w & 0 & 0 & x \\ 0 & 1 & w & y \\ 0 & 0 & 1 & z \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, D_z, D_w + xD_x + zD_y$.

- $A_{4.4}$ $[e_1, e_4] = e_1$, $[e_2, e_4] = e_1 + e_2$, $[e_3, e_4] = e_2 + e_3$:

$$S = \begin{bmatrix} e^w & we^w & \frac{w^2}{2}e^w & x \\ 0 & e^w & we^w & y \\ 0 & 0 & e^w & z \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, D_z, D_w + (x + y)D_x + (y + z)D_y + zD_z$.

- $A_{4.5ab}$ ($0 \leq ab$, $-1 \leq a \leq b \leq 1$) $[e_1, e_4] = e_1$, $[e_2, e_4] = ae_2$, $[e_3, e_4] = be_3$:

$$S = \begin{bmatrix} e^w & 0 & 0 & x \\ 0 & e^{aw} & 0 & y \\ 0 & 0 & e^{bw} & z \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, D_z, D_w + xD_x + ayD_y + bzD_z$.

- $A_{4.6ab}$ ($a \neq 0, b \geq 0$) $[e_1, e_4] = ae_1$, $[e_2, e_4] = be_2 - e_3$, $[e_3, e_4] = e_2 + be_3$:

$$S = \begin{bmatrix} e^{aw} & 0 & 0 & x \\ 0 & e^{bw} \cos w & e^{bw} \sin w & y \\ 0 & -e^{bw} \sin w & e^{bw} \cos w & z \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, D_z, D_w + axD_x + (by + z)D_y + (bz - y)D_z$.

- $A_{4.7}$ $[e_2, e_3] = e_1$, $[e_1, e_4] = 2e_1$, $[e_2, e_4] = e_2$, $[e_3, e_4] = e_2 + e_3$:

$$S = \begin{bmatrix} e^{2w} & -ze^w & (y - zw)e^w & x \\ 0 & e^w & we^w & y \\ 0 & 0 & e^w & z \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $-\frac{1}{2}D_x, zD_x + D_y, D_z - yD_x, D_w + 2xD_x + (y + z)D_y + zD_z$.

- $A_{4.8}, A_{4.9b}$ ($-1 \leq b \leq 1$) $[e_2, e_3] = e_1$, $[e_1, e_4] = (b + 1)e_1$, $[e_2, e_4] = e_2$, $[e_3, e_4] = be_3$:

$$S = \begin{bmatrix} e^{(b+1)w} & ye^{bw} & x \\ 0 & e^{bw} & z \\ 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, zD_x + D_y, -D_z, D_w + (xb + x)D_x + yD_y + bzD_z$.

- $A_{4.10}, A_{4.11}$ ($a \geq 0$) $[e_2, e_3] = e_1$, $[e_1, e_4] = 2ae_1$, $[e_2, e_4] = ae_2 - e_3$, $[e_3, e_4] = e_2 + ae_3$:

$$S = \begin{bmatrix} e^{2aw} & -e^{aw}(x \sin w + y \cos w) & e^{aw}(x \cos w - y \sin w) & z \\ 0 & e^{aw} \cos w & e^{aw} \sin w & x \\ 0 & -e^{aw} \sin w & e^{aw} \cos w & y \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $-2D_z, D_x + yD_z, D_y - xD_z, D_w + (ax + y)D_x + (ay - x)D_y + 2azD_z$.

- $A_{4.12}$ $[e_1, e_3] = e_1$, $[e_2, e_3] = e_2$, $[e_1, e_4] = -e_2$, $[e_2, e_4] = e_1$:

$$S = \begin{bmatrix} e^z \cos w & e^z \sin w & x \\ -e^z \sin w & e^z \cos w & y \\ 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, D_z + xD_x + yD_y, D_w + yD_x - xD_y$.

- $A_{5.1}$ $[e_3, e_5] = e_1$, $[e_4, e_5] = e_2$:

$$S = \begin{bmatrix} 1 & q & x & z \\ 0 & 1 & w & y \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $-D_q, -D_x, D_y, D_z, D_w - yD_q - zD_x$.

- $A_{5.2}$ $[e_2, e_5] = e_1$, $[e_3, e_5] = e_2$, $[e_4, e_5] = e_3$:

$$S = \begin{bmatrix} 1 & w & \frac{w^2}{2} & \frac{w^3}{6} & q \\ 0 & 1 & w & \frac{w^2}{2} & x \\ 0 & 0 & 1 & w & y \\ 0 & 0 & 0 & 1 & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_q, D_x, D_y, D_z, D_w + xD_q + yD_x + zD_y$.

- $A_{5.3}$ $[e_3, e_4] = e_2$, $[e_3, e_5] = e_1$, $[e_4, e_5] = e_3$:

$$S = \begin{bmatrix} 1 & 0 & -z & y - zw & q \\ 0 & 1 & w & \frac{w^2}{2} & x \\ 0 & 0 & 1 & w & 2y \\ 0 & 0 & 0 & 1 & 2z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $2D_x$, $-4D_q$, $D_y + 2zD_q$, $D_z - 2yD_q$, $D_w + 2yD_x + zD_y$.

- $A_{5.4}$ $[e_2, e_4] = e_1$, $[e_3, e_5] = e_1$:

$$S = \begin{bmatrix} 1 & x & y & q \\ 0 & 1 & 0 & z \\ 0 & 0 & 1 & w \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: D_q , D_z , D_w , $D_x + zD_q$, $D_y + wD_q$.

- $A_{5.5}$ $[e_3, e_4] = e_1$, $[e_2, e_5] = e_1$, $[e_3, e_5] = e_2$:

$$S = \begin{bmatrix} 1 & q & w + \frac{q^2}{2} & x \\ 0 & 1 & q & y \\ 0 & 0 & 1 & z \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: D_x , D_y , D_z , $D_w + zD_x$, $D_q + yD_x + zD_y$.

- $A_{5.6}$ $[e_3, e_4] = e_1$, $[e_2, e_5] = e_1$, $[e_3, e_5] = e_2$, $[e_4, e_5] = e_3$:

$$S = \begin{bmatrix} 1 & 2w & w^2 - z & y - zw + \frac{w^3}{3} & q \\ 0 & 1 & w & \frac{w^2}{2} & x \\ 0 & 0 & 1 & w & y \\ 0 & 0 & 0 & 1 & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $2D_q, -D_x, D_y + zD_q, -D_z + yD_q, -(D_w + 2xD_q + yD_x + zD_y)$.

- $A_{5.7abc}$ ($abc \neq 0, -1 \leq c \leq b \leq a \leq 1$) $[e_1, e_5] = e_1, [e_2, e_5] = ae_2, [e_3, e_5] = be_3, [e_4, e_5] = ce_4$:

$$S = \begin{bmatrix} e^w & 0 & 0 & 0 & q \\ 0 & e^{aw} & 0 & 0 & x \\ 0 & 0 & e^{bw} & 0 & y \\ 0 & 0 & 0 & e^{cw} & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_q, D_x, D_y, D_z, D_w + qD_q + axD_x + byD_y + czD_z$.

$$S = \begin{bmatrix} e^{aq} & 0 & we^{(a-1)q} & x \\ 0 & e^{bq} & ye^{(a-1)q} & z \\ 0 & 0 & e^{(a-1)q} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad (c = b - a + 1).$$

Right-invariant vector fields: $D_w, D_x, D_z, D_y, D_q + axD_x + (b + 1 - a)yD_y + bzD_z + wD_w$.

- $A_{5.8c}$ ($0 < c \leq 1$) $[e_2, e_5] = e_1, [e_3, e_5] = e_3, [e_4, e_5] = ce_4$:

$$S = \begin{bmatrix} e^{cw} & 0 & 0 & 0 & q \\ 0 & e^w & 0 & 0 & x \\ 0 & 0 & 1 & w & y \\ 0 & 0 & 0 & 1 & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, D_z, D_q, D_w + cqD_q + yD_x + zD_z$.

- $A_{5.9bc}$ ($0 \neq c \leq b$) $[e_1, e_5] = e_1, [e_2, e_5] = e_1 + e_2, [e_3, e_5] = be_3, [e_4, e_5] = ce_4$.

$$S = \begin{bmatrix} e^{cw} & 0 & 0 & 0 & q \\ 0 & e^{bw} & 0 & 0 & x \\ 0 & 0 & e^w & we^w & y \\ 0 & 0 & 0 & e^w & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_y, D_z, D_x, D_q, D_w + bxD_x + cqD_q + (y+z)D_y + zD_z$.

- $A_{5.10}$ $[e_2, e_5] = e_1, [e_3, e_5] = e_2, [e_4, e_5] = e_4$:

$$S = \begin{bmatrix} e^w & 0 & 0 & 0 & q \\ 0 & 1 & w & \frac{w^2}{2} & x \\ 0 & 0 & 1 & w & y \\ 0 & 0 & 0 & 1 & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right invariant vector fields: $D_x, D_y, D_z, D_q, D_w + qD_q + yD_x + zD_y$.

- $A_{5.11c}$ ($c \neq 0$) $[e_1, e_5] = e_1, [e_2, e_5] = e_1 + e_2, [e_3, e_5] = e_2 + e_3, [e_4, e_5] = ce_4$:

$$S = \begin{bmatrix} e^{cw} & 0 & 0 & 0 & q \\ 0 & e^w & we^w & \frac{w^2 e^w}{2} & x \\ 0 & 0 & e^w & we^w & y \\ 0 & 0 & 0 & e^w & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, D_z, D_q, D_w + (x+y)D_x + (y+z)D_y + zD_z + cqD_q$.

$$S = \begin{bmatrix} e^q & qe^q & w & x \\ 0 & e^q & y & z \\ 0 & 0 & 1 & q \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad (c = 1).$$

Right invariant vector fields: $D_x, \frac{1}{2}(D_z - D_w - qD_x), -\frac{1}{2}(D_y + qD_z), \frac{1}{2}(D_w + D_z + qD_x), D_q + (z+x)D_x + yD_y + zD_z + (y+w)D_w$.

- $A_{5.12}$ $[e_1, e_5] = e_1, [e_2, e_5] = e_1 + e_2, [e_3, e_5] = e_2 + e_3, [e_4, e_5] = e_3 + e_4$.

$$S = \begin{bmatrix} e^w & we^w & \frac{w^2 e^w}{2} & \frac{w^3 e^w}{6} & q \\ 0 & e^w & we^w & \frac{w^2 e^w}{2} & x \\ 0 & 0 & e^w & we^w & y \\ 0 & 0 & 0 & e^w & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_q, D_x, D_y, D_z, D_w + (q+x)D_q + (x+y)D_x + (y+z)D_y + zD_z + qD_q$.

- $A_{5.13apr}$ ($r \neq 0, 0 < |a| \leq 1$) $[e_1, e_5] = e_1, [e_2, e_5] = ae_2, [e_3, e_5] = pe_3 - re_4, [e_4, e_5] = re_3 + pe_4$:

$$S = \begin{bmatrix} e^w & 0 & 0 & 0 & q \\ 0 & e^{aw} & 0 & 0 & x \\ 0 & 0 & e^{pw} \cos rw & e^{pw} \sin rw & y \\ 0 & 0 & -e^{pw} \sin rw & e^{pw} \cos rw & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_q, D_x, D_y, D_z, D_w + qD_q + axD_x + (py + rz)D_y + (pz - ry)D_z$.

$$S = \begin{bmatrix} e^{-q} \cos \frac{rq}{2} & e^{-q} \sin \frac{rq}{2} & w & x \\ -e^{-q} \sin \frac{rq}{2} & e^{-q} \cos \frac{rq}{2} & y & z \\ 0 & 0 & \cos \frac{rq}{2} & \sin \frac{rq}{2} \\ 0 & 0 & -\sin \frac{rq}{2} & \cos \frac{rq}{2} \end{bmatrix} \quad (a = p = 1).$$

Right-invariant vector fields:

$$\begin{aligned} & \frac{1}{2} \left(\cos \frac{rq}{2} D_x - \cos \frac{rq}{2} D_y - \sin \frac{rq}{2} D_z - \sin \frac{rq}{2} D_w \right), \\ & \frac{1}{2} \left(\left(\cos \frac{rq}{2} - \sin \frac{rq}{2} \right) D_w + \left(\cos \frac{rq}{2} + \sin \frac{rq}{2} \right) D_x \right. \\ & \quad \left. - \left(\cos \frac{rq}{2} + \sin \frac{rq}{2} \right) D_y + \left(\cos \frac{rq}{2} - \sin \frac{rq}{2} \right) D_z \right), \\ & \frac{1}{2} \left(\cos \frac{rq}{2} D_w + \sin \frac{rq}{2} D_x + \sin \frac{rq}{2} D_y - \cos \frac{rq}{2} D_z \right), \\ & \frac{1}{2} \left(\sin \frac{rq}{2} D_w - \cos \frac{rq}{2} D_y - \sin \frac{rq}{2} D_z - \cos \frac{rq}{2} D_x \right), \\ & \left(x - \frac{rz}{2} \right) D_x + \left(\frac{rw}{2} + y \right) D_y + \left(\frac{rx}{2} + z \right) D_z + \left(w - \frac{ry}{2} \right) D_w - D_q. \end{aligned}$$

- $A_{5.14p}$ $[e_2, e_5] = e_1, [e_3, e_5] = pe_3 - e_4, [e_4, e_5] = e_3 + pe_4$:

$$S = \begin{bmatrix} 1 & x & 0 & 0 & q \\ 0 & 1 & 0 & 0 & w \\ 0 & 0 & e^{pw} \cos w & e^{pw} \sin w & y \\ 0 & 0 & -e^{pw} \sin w & e^{pw} \cos w & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_q, D_x, D_y, D_z, D_w + xD_q + (py + z)D_y + (pz - y)D_z$.

- $A_{5.15a}$ $[e_1, e_5] = e_1, [e_2, e_5] = e_1 + e_2, [e_3, e_5] = ae_3, [e_4, e_5] = e_3 + ae_4$:

$$S = \begin{bmatrix} e^{aq} & qe^{aq} & we^{(a-1)q} & x \\ 0 & e^{aq} & ye^{(a-1)q} & z \\ 0 & 0 & e^{(a-1)q} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_w, D_y, D_x, D_z, D_q + (z + ax)D_x + yD_y + azD_z + (w + y)D_w$.

- $A_{5.16pr}$ ($r > 0$) $[e_1, e_5] = e_1, [e_2, e_5] = e_1 + e_2, [e_3, e_5] = pe_3 - re_4, [e_4, e_5] = re_3 + pe_4$:

$$S = \begin{bmatrix} e^w & we^w & 0 & 0 & q \\ 0 & e^w & 0 & 0 & x \\ 0 & 0 & e^{pw} \cos rw & e^{pw} \sin rw & y \\ 0 & 0 & -e^{pw} \sin rw & e^{pw} \cos rw & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_q, D_x, D_y, D_z, D_w + (q + x)D_q + xD_x + (py + rz)D_y + (pz - ry)D_z$.

- $A_{5.17prs}$ ($s > 0$) $[e_1, e_5] = pe_1 - e_2, [e_2, e_5] = e_1 + pe_2, [e_3, e_5] = re_3 - se_4, [e_4, e_5] = se_3 + re_4$:

$$S = \begin{bmatrix} e^{pw} \cos w & e^{pw} \sin w & 0 & 0 & x \\ -e^{pw} \sin w & e^{pw} \cos w & 0 & 0 & y \\ 0 & 0 & e^{rw} \cos sw & e^{rw} \sin sw & z \\ 0 & 0 & -e^{rw} \sin sw & e^{rw} \cos sw & q \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, D_z, D_q, D_w + (px + y)D_x + (py - x)D_y + (rz + sq)D_z + (rq - sz)D_q$.

$$S = \begin{bmatrix} e^{pq} \cos \frac{(s+1)q}{2} & e^{pq} \sin \frac{(s+1)q}{2} & w & x \\ -e^{pq} \sin \frac{(s+1)q}{2} & e^{pq} \cos \frac{(s+1)q}{2} & y & z \\ 0 & 0 & \cos \frac{(s-1)q}{2} & \sin \frac{(s-1)q}{2} \\ 0 & 0 & -\sin \frac{(s-1)q}{2} & \cos \frac{(s-1)q}{2} \end{bmatrix} \quad (r = p).$$

Right-invariant vector fields:

$$\begin{aligned} & \cos \frac{(s-1)q}{2} D_z - \sin \frac{(s-1)q}{2} D_y - \sin \frac{(s-1)q}{2} D_x - \cos \frac{(s-1)q}{2} D_w, \\ & \cos \frac{(s-1)q}{2} D_y + \cos \frac{(s-1)q}{2} D_x - \sin \frac{(s-1)q}{2} D_w + \sin \frac{(s-1)q}{2} D_z, \\ & \sin \frac{(s-1)q}{2} D_x - \sin \frac{(s-1)q}{2} D_y + \cos \frac{(s-1)q}{2} D_w + \cos \frac{(s-1)q}{2} D_z, \\ & -\cos \frac{(s-1)q}{2} D_y + \cos \frac{(s-1)q}{2} D_x - \sin \frac{(s-1)q}{2} D_w - \sin \frac{(s-1)q}{2} D_z, \\ & D_q + \frac{(s+1)z+2py}{2} D_y - \frac{(s+1)z-2px}{2} D_x - \frac{(s+1)y-2pw}{2} D_w + \frac{(s+1)x+2pz}{2} D_z. \end{aligned}$$

$$S = \begin{bmatrix} \cos q & \sin q & w & x \\ -\sin q & \cos q & y & z \\ 0 & 0 & e^{pq} & 0 \\ 0 & 0 & 0 & e^{rq} \end{bmatrix} \quad (s = 1).$$

Right-invariant vector fields: $e^{pq}D_w, e^{pq}D_y, e^{rq}D_x, e^{rq}, -D_q - zD_x + wD_y + xD_z - yD_w$.

- $A_{5,18p} (p \geq 0)$ $[e_1, e_5] = pe_1 - e_2, [e_2, e_5] = e_1 + pe_2, [e_3, e_5] = e_1 + pe_3 - e_4, [e_4, e_5] = e_2 + e_3 + pe_4$

$$S = \begin{bmatrix} \cos q & \sin q & w & x \\ -\sin q & \cos q & y & z \\ 0 & 0 & e^{pq} & qe^{pq} \\ 0 & 0 & 0 & e^{pq} \end{bmatrix}.$$

Right-invariant vector fields: $e^{pq}D_x, -e^{pq}D_z, e^{pq}(D_w + qD_x), -e^{pq}(D_y + qD_z), -D_q - zD_x + wD_y + xD_z - yD_w$.

- $A_{5.19ab}$ ($b \neq 0$) $[e_2, e_3] = e_1$, $[e_1, e_5] = ae_1$, $[e_2, e_5] = e_2$, $[e_3, e_5] = (a-1)e_3$, $[e_4, e_5] = be_4$:

$$S = \begin{bmatrix} e^{aw} & e^w x & 0 & z \\ 0 & e^w & 0 & y \\ 0 & 0 & e^{bw} & q \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: D_z , D_y , $D_x + yD_z$, D_q , $D_w + bqD_q + (a-1)xD_x + yD_y + azD_z$.

- $A_{5.20a}$ $[e_2, e_3] = e_1$, $[e_1, e_5] = ae_1$, $[e_2, e_5] = e_2$, $[e_3, e_5] = (a-1)e_3$, $[e_4, e_5] = e_1 + ae_4$:

$$S = \begin{bmatrix} e^{aw} & e^w x & we^{aw} & z \\ 0 & e^w & 0 & y \\ 0 & 0 & e^{aw} & q \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: D_z , D_y , $D_x + yD_z$, D_q , $D_w + aqD_q + (a-1)xD_x + yD_y + (q + az)D_z$.

- $A_{5.21}$ $[e_2, e_3] = e_1$, $[e_1, e_5] = 2e_1$, $[e_2, e_5] = e_2 + e_3$, $[e_3, e_5] = e_3 + e_4$, $[e_4, e_5] = e_4$:

$$S = \begin{bmatrix} e^{2w} & 0 & ze^w & (z - y + zw)e^w & q \\ 0 & e^w & we^w & \frac{w^2 e^w}{2} & x \\ 0 & 0 & e^w & we^w & y \\ 0 & 0 & 0 & e^w & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $-2D_q$, $(y + z)D_q + D_z$, $D_y - zD_q$, D_x , $D_w + 2qD_q + (x + y)D_x + (y + z)D_y + zD_z$.

- $A_{5.22}$ $[e_2, e_3] = e_1$, $[e_2, e_5] = e_3$, $[e_4, e_5] = e_4$:

$$S = \begin{bmatrix} e^w & 0 & 0 & 0 & q \\ 0 & 1 & z & \frac{z^2}{2} & x \\ 0 & 0 & 1 & z & y \\ 0 & 0 & 0 & 1 & w \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_z + yD_x + wD_y, -D_y, D_q, D_w + qD_q$.

- $A_{5,23b}$ ($b \neq 0$) $[e_2, e_3] = e_1, [e_1, e_5] = 2e_1, [e_2, e_5] = e_2 + e_3, [e_3, e_5] = e_3, [e_4, e_5] = be_4$:

$$S = \begin{bmatrix} e^{bw} & 0 & 0 & 0 & q \\ 0 & e^{2w} & -ze^w & ye^w & x \\ 0 & 0 & e^w & we^w & y + zw \\ 0 & 0 & 0 & e^w & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $-\frac{1}{2}D_x, zD_x + D_y, D_z - (y + zw)D_x - wD_y, D_q, D_w + qD_q + 2xD_x + yD_y + zD_z$.

$$S = \begin{bmatrix} 1 & ye^w & (q - yw - 2ze^w) & xe^{2w} \\ 0 & e^w & -we^w & qe^{2w} \\ 0 & 0 & e^w & ye^{2w} \\ 0 & 0 & 0 & e^{2w} \end{bmatrix} \quad (b = 1).$$

Right-invariant vector fields: $-2D_x, D_y + qD_x, D_z + D_q - yD_x, \frac{1}{2}(D_q + yD_x), -D_w + (q + y)D_q + 2xD_x + yD_y + (y + z)D_z$.

- $A_{5,24\epsilon}$ ($\epsilon = \pm 1$) $[e_2, e_3] = e_1, [e_1, e_5] = 2e_1, [e_2, e_5] = e_2 + e_3, [e_3, e_5] = e_3, [e_4, e_5] = \epsilon e_1 + 2e_4$:

$$S = \begin{bmatrix} 1 & q & ze^q & xe^{2q} \\ 0 & 1 & ye^q & \left(\frac{y^2}{2} - \epsilon w\right)e^{2q} \\ 0 & 0 & e^q & ye^{2q} \\ 0 & 0 & 0 & e^{2q} \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, -D_z - yD_x, D_w, -D_q + (2x - \frac{y^2}{2} + \epsilon w)D_x + yD_y + (z - y)D_z + 2wD_w$.

- $A_{5,25bp}$ ($b \neq 0$) $[e_2, e_3] = e_1, [e_1, e_5] = 2pe_1, [e_2, e_5] = pe_2 + e_3, [e_3, e_5] = pe_3 - e_2, [e_4, e_5] = be_4$:

$$S = \begin{bmatrix} e^{2pw} & -e^{pw}(y \cos w + x \sin w) & e^{pw}(x \cos w - y \sin w) & 0 & z \\ 0 & e^{pw} \cos w & e^{pw} \sin w & 0 & x \\ 0 & -e^{pw} \sin w & e^{pw} \cos w & 0 & y \\ 0 & 0 & 0 & e^{bw} & q \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $2D_z, D_x + yD_z, -D_y + xD_z, D_q, D_w + bqD_q + (px + y)D_x + (py - x)D_y + 2pzD_z$.

- $A_{5.26\epsilon p}$ ($\epsilon = \pm 1$) $[e_2, e_3] = e_1, [e_1, e_5] = 2pe_1, [e_2, e_5] = pe_2 + e_3, [e_3, e_5] = pe_3 - e_2, [e_4, e_5] = \epsilon e_1 + 2pe_4$:

$$S = \begin{bmatrix} e^{2pw} & -e^{pw}(y \cos w + x \sin w) & e^{pw}(x \cos w - y \sin w) & 2\epsilon we^{2pw} & z \\ 0 & e^{pw} \cos w & e^{pw} \sin w & 0 & x \\ 0 & -e^{pw} \sin w & e^{pw} \cos w & 0 & y \\ 0 & 0 & 0 & e^{2pw} & q \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $2D_z, D_x + yD_z, -D_y + xD_z, D_q, D_w + (px + y)D_x + (py - x)D_y + 2(\epsilon q + pz)D_z + 2pqD_q$.

- $A_{5.27}$ $[e_2, e_3] = e_1, [e_1, e_5] = e_1, [e_3, e_5] = e_3 + e_4, [e_4, e_5] = e_1 + e_4$:

$$S = \begin{bmatrix} e^w & we^w & \left(q + \frac{w^2}{2}\right)e^w & z \\ 0 & e^w & we^w & x \\ 0 & 0 & e^w & y \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_z, -D_q - yD_z, D_y, D_x, D_w + (x + y)D_x + yD_y + (x + z)D_z$.

- $A_{5.28a}$ $[e_2, e_3] = e_1, [e_1, e_5] = ae_1, [e_2, e_5] = (a - 1)e_2, [e_3, e_5] = e_3 + e_4, [e_4, e_5] = e_4$:

$$S = \begin{bmatrix} e^{-w} & z & xe^{-(a-1)w} & q \\ 0 & 1 & ye^{(a-1)w} & w \\ 0 & 0 & e^{(a-1)w} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $e^{2(a-1)w}D_x, D_y, ye^{(2(a-1)w)}D_x + D_z + wD_q, D_q, -D_w - (a-2)xD_x + (a-1)yD_y + zD_z + qD_q$.

- $A_{5.29}$ $[e_2, e_4] = e_1, [e_1, e_5] = e_1, [e_2, e_5] = e_2, [e_4, e_5] = e_3$:

$$S = \begin{bmatrix} 1 & q & xe^w & z \\ 0 & 1 & ye^w & w \\ 0 & 0 & e^w & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_x, D_y, D_z, D_q + yD_x + wD_z, -D_w + xD_x + yD_y$.

- $A_{5.30a}$ $[e_2, e_4] = e_1, [e_3, e_4] = e_2, [e_1, e_5] = (a+1)e_1, [e_2, e_5] = ae_2, [e_3, e_5] = (a-1)e_3, [e_4, e_5] = e_4$:

$$S = \begin{bmatrix} 1 & q & \frac{q^2}{2} & x \\ 0 & e^w & e^w q & e^w y \\ 0 & 0 & e^{2w} & e^{2w} z \\ 0 & 0 & 0 & e^{(a+1)w} \end{bmatrix}.$$

Right-invariant vector fields: $e^{(a+1)w}D_x, e^{aw}D_y, e^{(a-1)w}D_z, e^w(D_q + yD_x + zD_y), -D_w$.

- $A_{5.31}$ $[e_2, e_4] = e_1, [e_3, e_4] = e_2, [e_1, e_5] = 3e_1, [e_2, e_5] = 2e_2, [e_3, e_5] = e_3, [e_4, e_5] = e_3 + e_4$:

$$S = \begin{bmatrix} e^{3w} & -ze^{2w} & \frac{1}{2}z^2e^w & \frac{1}{2}e^w(z^2w + x - yz + \frac{3z^2}{2}) & q \\ 0 & e^{2w} & -ze^w & e^w(y - z - zw) & x \\ 0 & 0 & e^w & we^w & y \\ 0 & 0 & 0 & e^w & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $3D_q, -(2D_x + zD_q), D_y + zD_x, D_z - xD_q - (y+z)D_x, D_w + 3qD_q + 2xD_x + (y+z)D_y + zD_z$.

- $A_{5.32a}$ $[e_2, e_4] = e_1, [e_3, e_4] = e_2, [e_1, e_5] = e_1, [e_2, e_5] = e_2, [e_3, e_5] = ae_1 + e_3$:

$$S = \begin{bmatrix} 1 & q & \frac{q^2}{2} - aw & x \\ 0 & 1 & q & y \\ 0 & 0 & 1 & z \\ 0 & 0 & 0 & e^w \end{bmatrix}.$$

Right-invariant vector fields: $e^w D_x$, $e^w D_y$, $e^w D_z$, $D_q + yD_x + zD_y$, $-D_w + azD_x$.

- $A_{5.33ab}$ ($a^2 + b^2 \neq 0$) $[e_1, e_4] = e_1$, $[e_3, e_4] = be_3$, $[e_2, e_5] = e_2$, $[e_3, e_5] = ae_3$:

$$S = \begin{bmatrix} e^z & 0 & 0 & q \\ 0 & e^w & 0 & x \\ 0 & 0 & e^{aw+bz} & y \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: D_q , D_x , D_y , $-(D_z + qD_q + byD_y)$, $-(D_w + xD_x + ayD_y)$.

- $A_{5.34a}$ $[e_1, e_4] = ae_1$, $[e_2, e_4] = e_2$, $[e_3, e_4] = e_3$, $[e_1, e_5] = e_1$, $[e_3, e_5] = e_2$.

$$S = \begin{bmatrix} e^{\alpha z+w} & 0 & 0 & q \\ 0 & e^z & w & x \\ 0 & 0 & e^z & y \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: D_q , D_x , D_y , $D_z + xD_x + yD_y + aqD_q$, $D_w + yD_x + qD_q$.

- $A_{5.35ab}$ ($a^2 + b^2 \neq 0$) $[e_1, e_4] = be_1$, $[e_2, e_4] = e_2$, $[e_3, e_4] = e_3$, $[e_1, e_5] = ae_1$, $[e_2, e_5] = -e_3$, $[e_3, e_5] = e_2$:

$$S = \begin{bmatrix} e^{aw+bz} & 0 & 0 & q \\ 0 & e^z \cos w & e^z \sin w & x \\ 0 & -e^z \sin w & e^z \cos w & y \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: D_q , D_x , D_y , $D_z + xD_x + yD_y + bqD_q$, $D_w y D_x - x D_y + aq D_q$.

- $A_{5.36}$ $[e_2, e_3] = e_1, [e_1, e_4] = e_1, [e_2, e_4] = e_2, [e_2, e_5] = -e_2, [e_3, e_5] = e_3$:

$$S = \begin{bmatrix} e^w & e^q x & z \\ 0 & e^q & y \\ 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_z, -D_x - yD_z, D_y, Dw + xD_x + zD_z, D_q - xD_x + yD_y$.

- $A_{5.37}$ $[e_2, e_3] = e_1, [e_1, e_4] = 2e_1, [e_2, e_4] = e_2, [e_3, e_4] = e_3, [e_2, e_5] = -e_3, [e_3, e_5] = e_2$:

$$S = \begin{bmatrix} e^{2q} & (y \cos w + x \sin w)e^q & (y \sin w - x \cos w)e^q & z \\ 0 & e^q \cos w & e^q \sin w & x \\ 0 & -e^q \sin w & e^q \cos w & y \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $2D_z, D_x - yD_z, D_y + xD_z, D_q + xD_x + yD_y + 2zD_z, D_w + yD_x - xD_y$.

- $A_{5.38}$ $[e_1, e_4] = e_1, [e_2, e_5] = e_2, [e_4, e_5] = e_3$:

$$S = \begin{bmatrix} e^z & 0 & 0 & 0 & q \\ 0 & e^w & 0 & 0 & x \\ 0 & 0 & 1 & w & y \\ 0 & 0 & 0 & 1 & z \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_q, D_x, D_y, qD_q + D_z, D_w + xD_x + zD_y$.

- $A_{5.39}$ $[e_1, e_4] = e_1, [e_2, e_4] = e_2, [e_1, e_5] = -e_2, [e_2, e_5] = e_1, [e_4, e_5] = e_3$:

$$S = \begin{bmatrix} 1 & z & 0 & 0 & q \\ 0 & 1 & 0 & 0 & w \\ 0 & 0 & e^{-w} \cos z & e^{-w} \sin z & x \\ 0 & 0 & -e^{-w} \sin z & e^{-w} \cos z & y \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $D_y, D_x, D_q, -D_w + yD_y + xD_x, -D_z + xD_y - yD_x - wD_q$.

- $A_{5,40}$ $[e_1, e_2] = 2e_1, [e_3, e_1] = e_2, [e_2, e_3] = 2e_3, [e_1, e_4] = e_5, [e_2, e_4] = e_4, [e_2, e_5] = -e_5, [e_3, e_5] = e_4$:

$$S = \begin{bmatrix} e^x & y & w \\ z & (1 + yz)e^{-x} & q \\ 0 & 0 & 1 \end{bmatrix}.$$

Right-invariant vector fields: $ze^{-x}D_x + (yz + 1)e^{-x}D_y + qD_w, -qD_q + D_x + yD_y - zD_z + wD_w, wD_q + e^xD_z, D_q, D_w$.

9. APPENDIX

PROPOSITION 9.1. *Algebra $A_{5,21}$ has no representation in $\mathfrak{gl}(4, \mathbb{R})$.*

Proof. First of all assume that there is a upper triangular representation. Since each of e_1, e_2, e_3, e_4 is a sum of commutators we may assume that each of E_1, E_2, E_3, E_4 are strictly upper triangular. Now put

$$E_2 = \begin{bmatrix} 0 & t & u & v \\ 0 & 0 & p & q \\ 0 & 0 & 0 & r \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad E_5 = \begin{bmatrix} a & b & c & d \\ 0 & e & f & g \\ 0 & 0 & h & i \\ 0 & 0 & 0 & j \end{bmatrix},$$

and define $E_3 = [E_2, E_5] - E_2, E_4 = [E_3, E_5] - E_3, E_1 = [E_2, E_3]$. Then

$$E_1 = \begin{bmatrix} 0 & 0 & tp(h - 2e + a) & tpi + tqj - 2teq - 2tfr + urj - 2urh + tqa + rau + rbp \\ 0 & 0 & 0 & pr(j - 2h + e) \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$E_3 = \begin{bmatrix} 0 & t(e - a - 1) & tf + uh - au - bp - u & tg + ui + vj - av - bq - cr - v \\ 0 & 0 & p(h - e - 1) & pi + qj - eq - fr - q \\ 0 & 0 & 0 & r(j - h - 1) \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$E_4 = \begin{bmatrix} 0 & t(a-e+1)^2 & * & * \\ 0 & 0 & p(-h+e+1)^2 & * \\ 0 & 0 & 0 & r(-j+h+1)^2 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

It turns out that $[E_1, E_5] = [E_2, E_4]$, the equality of commutators. Furthermore we have that $[E_2, E_5] = E_2 + E_3$, $[E_3, E_5] = E_3 + E_4$, $[E_2, E_3] = E_1$ by construction. The remaining brackets are given by

$$[E_1, E_2] = \begin{bmatrix} 0 & 0 & 0 & tpr(3h-3e+a-j) \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$[E_1, E_3] = \begin{bmatrix} 0 & 0 & 0 & tpr(jh-h^2-3h-3je+4he+3e+2ja-3ha-a-e^2+ea+j) \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$[E_1, E_4] = \begin{bmatrix} 0 & 0 & 0 & * \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$[E_1, E_5] - 2E_5 = \begin{bmatrix} 0 & 0 & tp(h-a-2)(h-2e+a) & * \\ 0 & 0 & 0 & pr(j-e-2)(j-2h+e) \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$[E_1, E_4] = \begin{bmatrix} 0 & 0 & 0 & * \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$[E_3, E_4] = \begin{bmatrix} 0 & 0 & tp(h-e-1)(e-a-1)(h-2e+a) & * \\ 0 & 0 & 0 & pr(j-h-1)(h-e-1)(j-2h+e) \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$[E_4, E_5] - E_4 = \begin{bmatrix} 0 & t(e-a-1)^3 & * & * \\ 0 & 0 & p(h-e-1)^3 & * \\ 0 & 0 & 0 & r(j-h-1)^3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Looking at the $[E_1, E_2]$ bracket we see that $prt(3h - 3e + a - j) = 0$. Suppose first of all that $r = 0$. Then $t \neq 0$ or else $E_1 = 0$. From the $(1, 2)$ -entry in the $[E_4, E_5]$ bracket we deduce that $e = a + 1$. Now it must be that $p \neq 0$ or else $[E_1, E_5] = 2E_5$ implies that $E_1 = 0$. Now from the $(1, 3)$ -entry in the $[E_1, E_5]$ bracket we deduce that $h = a + 2$. From the $(1, 3)$ -entry in the $[E_4, E_5]$ bracket we conclude that $u = bp - ft$. However, we now have a contradiction because E_1 and E_4 are proportional.

Hence we may assume that $r \neq 0$ and by appealing to Corollary 3.2 that also $t \neq 0$ and hence $rt \neq 0$. Now suppose that $p = 0$. Then comparing E_1 and the $[E_1, E_5]$ bracket we find that $j = a + 2$. From the $(1, 2)$ and $(4, 5)$ -entries in the $[E_4, E_5]$ bracket we find that $e = a + 1$, $h = a + 1$ which is a contradiction because now E_1 and E_4 are proportional.

Hence we assume that $prt \neq 0$. Then from the $(1, 2)$, $(2, 3)$ and $(4, 5)$ -entries in the $[E_4, E_5]$ bracket we find that $e = a + 1$, $h = e + 1$, $j = h + 1$ which is a contradiction because now E_1, E_3 and E_4 are linearly dependent. Hence there can be no representation of $A_{5,21}$ in $\mathfrak{gl}(4, \mathbb{R})$. ■

PROPOSITION 9.2. *For algebra $A_{5,22}$ we have $\mu = 5$.*

We assume that algebra $A_{5,22}$ has an upper triangular representation. In fact we may assume that E_2 and E_5 are upper triangular and that E_4 is strictly upper triangular, E_1 and E_3 being determined by the brackets that define the algebra. We shall write

$$E_2 = \begin{bmatrix} \alpha & \beta & \delta & \rho \\ \alpha & \beta & \delta & \rho \\ 0 & \lambda & \sigma & \tau \\ 0 & 0 & \phi & \mu \end{bmatrix}, \quad E_5 = \begin{bmatrix} c & d & e & f \\ 0 & g & h & i \\ 0 & 0 & j & k \\ 0 & 0 & 0 & m \end{bmatrix}.$$

From the (1, 2), (2, 3), (3, 4)-entries of $[E_1, E_2] = 0$ it follows that the (1, 2), (2, 3), (3, 4)-entries of E_1 are zero. From $[E_2, E_3] = E_1$ and $[E_3, E_5] = 0$ it follows that the (1, 2), (2, 3), (3, 4)-entries of E_3 are zero. Invoking Corollary 3.2 we make an upper-triangular transformation so as to reduce E_4 to one of the following seven forms and work through each of these seven cases in turn:

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$\begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

In the interests of saving space we do not write out explicitly the very complicated matrices concerned. In the first case from $[E_2, E_4] = 0$ we find that $\phi = \alpha$, $\sigma = \beta$, $\tau = \delta$, $\lambda = \alpha$, $\mu = \beta$, $\nu = \alpha$ and from $[E_4, E_5] = E_4$ that $g = c + 1$, $h = d$, $i = e$, $j = g + 1$, $k = h$, $t = j + 1$. Now, however, $[E_3, E_5] = 0$ implies that $E_3 = 0$.

In the second case from $[E_2, E_4] = 0$ we find that $\phi = \alpha$, $\sigma = \beta$, $\lambda = \alpha$, $\tau = 0$, $\mu = 0$ and from $[E_4, E_5] = E_4$ that $g = c + 1$, $h = d$, $i = 0$, $j = g + 1$, $k = 0$. Now, however, $[E_1, E_2] = 0$ implies that $E_1 = 0$.

In the third case from $[E_2, E_4] = 0$ we find that $\phi = \alpha$, $\sigma = 0$, $\tau = \delta$, $\nu = \lambda$ and from $[E_4, E_5] = E_4$ that $g = c + 1$, $h = 0$, $i = e$, $t = j + 1$. The (1, 3) entries of $[E_1, E_2] = 0$ $[E_3, E_5] = 0$ imply that $e\alpha + j\delta - c\delta - e\nu = 0$ and hence E_1 and E_3 are proportional.

In the fourth case from $[E_2, E_4] = 0$ we find that $\lambda = \alpha$, $\mu = \beta$, $\nu = \phi$ and from $[E_4, E_5] = E_4$ that $j = c + 1$, $k = d$, $t = g + 1$. Now the only non-zero entry of the commutator $[E_2, E_3]$ is in the (1, 4) position. On the hand the (1, 3) and (2, 4) entries of $[E_3, E_5]$ imply that $h\beta + \delta - d\sigma = 0$ and $d\sigma + \tau - h\beta = 0$ and hence the only non-zero entry of E_3 is in the (1, 4) position. Thus if also $[E_2, E_3] = E_1$ then E_1 and E_3 would be proportional.

In the fifth case from $[E_2, E_4] = 0$ we find that $\lambda = \alpha$, $\mu = 0$ and from $[E_4, E_5] = E_4$ that $j = c + 1$, $k = 0$. The (1, 3)-entry of $[E_3, E_5] = 0$ implies that the (1, 3)-entry of E_3 is zero and (2, 4)-entries of $[E_1, E_2] = 0$ and $[E_3, E_5] = 0$ that the (2, 4)-entry of E_3 is zero. Now $[E_2, E_3] = E_1$ gives that E_1 and E_3 are proportional.

In the sixth case from $[E_2, E_4] = 0$ we find that $\phi = \lambda$, $\beta = 0$, $\mu = 0$ and from $[E_4, E_5] = E_4$ that $d = 0$, $j = g + 1$, $k = 0$. Now, however, $[E_1, E_2] = 0$ immediately implies that $E_1 = 0$.

In the seventh case from $[E_2, E_4] = 0$ we find that $\nu = \alpha$ and from $[E_4, E_5] = E_4$ that $t = c + 1$. Then $[E_2, E_3] = E_1$ implies that the $(1, 3)$ entry of E_1 is zero. If $(\alpha - \lambda)(\alpha - \phi) \neq 0$ then comparing E_1 and $[E_1, E_2] = 0$ implies that E_1 and E_4 are proportional. Hence by appealing to Corollary 3.2 we may assume that $\lambda = \alpha$. Again comparing E_1 and $[E_1, E_2]$ we find that $\phi = \alpha$ in order not to have E_1 and E_4 proportional. Now, however, $[E_2, E_3] = E_1$ gives that E_1 and E_4 are proportional.

PROPOSITION 9.3. $\mu(\mathfrak{g}) = 5$ for algebra $A_{5,23}^b$ for $b \neq 1$ and $\mu(\mathfrak{g}) = 4$ for $b = 1$.

We define

$$E_2 = \begin{bmatrix} 0 & \beta & \delta & \rho \\ 0 & 0 & \sigma & \tau \\ 0 & 0 & 0 & \mu \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad E_4 = \begin{bmatrix} 0 & m & n & p \\ 0 & 0 & q & r \\ 0 & 0 & 0 & s \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad E_5 = \begin{bmatrix} c & d & e & f \\ 0 & g & h & i \\ 0 & 0 & j & k \\ 0 & 0 & 0 & t \end{bmatrix},$$

and then E_3 is defined as $[E_2, E_5] - E_2$ and E_1 as $[E_2, E_3]$ giving

$$E_3 = \begin{bmatrix} 0 & -\beta(-g+c+1) & \beta h + \delta j - c\delta - d\sigma - \delta & \beta i + \delta k + \rho t - c\rho - d\tau - e\mu - \rho \\ 0 & 0 & -\sigma(-j+g+1) & \sigma k + \tau t - g\tau - h\mu - \tau \\ 0 & 0 & 0 & -\mu(-t+j+1) \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$E_1 = \begin{bmatrix} 0 & 0 & \beta\sigma(j-2g+c) & \beta\sigma k + \beta\tau t - 2\beta g\tau - 2\beta h\mu + \delta\mu t - 2\delta\mu j + \beta\tau c + \mu c\delta + \mu d\sigma \\ 0 & 0 & 0 & \sigma\mu(t-2j+g) \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

To begin with we consider the $(1, 2)$ -entry of $[E_3, E_5] = E_3$ which gives $\beta(g - c - 1) = 0$. We separate cases according as first of all $\beta = 0$ and then $g - c - 1 = 0$. Looking at the $(3, 4)$ -entry of $[E_3, E_5] = E_3$ we must have that $\mu \neq 0$ or $E_1 = 0$ and so $t = j + 1$. Next, from $[E_4, E_5] = bE_4$, assuming that $b \neq 1$ we find that $s = 0$. From the $(2, 4)$ -entry of $[E_2, E_4] = 0$ we find that $q = 0$. Now comparing E_1 and $[E_1, E_5] - 2E_1$ we find that $j = c + 1$ in order

not to have $E_1 = 0$. Next from the (1, 3) and (1, 4)-entries of $[E_2, E_4] = 0$, respectively, we have $\sigma \neq 0$ or $\mu \neq 0$ or else $E_1 = 0$. From the (2, 3), (2, 4) and (1, 4)-entries of $[E_3, E_5] = E_3$, respectively, we have that $g = c$, $\tau = h\mu - \sigma k$ and $\rho = dh\mu - \delta k + e\mu$. Looking at the (2, 4)-entry of $[E_4, E_5] = bE_4$ we have $b = 2$ and $r \neq 0$ or else E_1 and E_4 are proportional. Now from the (1, 4)-entry of $[E_4, E_5] = bE_4$ we have that $d = 0$ and hence $E_3 = 0$.

Now we consider the second case where $g = c + 1$ and $\beta \neq 0$. From the (1, 2)-entry of $[E_4, E_5] - bE_4 = 0$ we find that $m = 0$ and from the (1, 3)-entry of $[E_2, E_4] = 0$ we find that $q = 0$. Comparing E_1 and $[E_1, E_5] - 2E_1$ we find that $t = c + 2$ in order not to have $E_1 = 0$. Now not both μ and σ can be zero or else $E_1 = 0$ so we distinguish subcases according as σ and μ zero. So if $\sigma = 0$ and $\mu \neq 0$ the (1, 2) and (3, 4)-entries of $[E_4, E_5] = bE_4$, respectively, give that $n = s = 0$, since we are assuming that $b \neq 1$. Finally the (1, 4)-entry of $[E_2, E_4]$ implies that r and hence E_1 and E_4 are proportional. On the other hand if $\mu = 0$ and $\sigma \neq 0$ then from the (1, 3)-entry of $[E_2, E_3] - E_1 = 0$ we find that $j = c + 2$ and from the (2, 4)-entry of $[E_2, E_4] = 0$ that $s = 0$. Next the (2, 4)-entry of $[E_4, E_5] - bE_4 = 0$ that $r = 0$, from the (1, 3)-entry of $[E_3, E_5] - E_3 = 0$ that $\delta = d\sigma - \beta h$ from the (1, 4)-entry of $[E_3, E_5] - E_3 = 0$ that $\rho = (k - i)\beta h + d\tau$. From the (1, 3)-entry of $[E_4, E_5] - bE_4 = 0$ we find that $b = 2$ and $n \neq 0$ or else E_1 and E_4 are proportional. Finally the (1, 4)-entry of $[E_4, E_5] - bE_4 = 0$ implies that $k = 0$ which gives that $E_1 = 0$.

PROPOSITION 9.4. Algebra $A_{5,31}$ has no representation in $\mathfrak{gl}(4, \mathbb{R})$.

Proof. Put

$$E_4 = \begin{bmatrix} 0 & t & u & v \\ 0 & 0 & p & q \\ 0 & 0 & 0 & r \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad E_5 = \begin{bmatrix} a & b & c & d \\ 0 & e & f & g \\ 0 & 0 & h & i \\ 0 & 0 & 0 & j \end{bmatrix},$$

and define $E_3 = [E_4, E_5] - E_4$, $E_2 = [E_3, E_4]$, $E_1 = [E_2, E_4]$. Then

$$E_1 = \begin{bmatrix} 0 & 0 & 0 & tpr(3e - a - 3h + j) \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$E_2 = \begin{bmatrix} 0 & 0 & tp(2e - a - h) & 2tqe - tqa + 2rtf + 2ruh - rau - rbp - tpi - tqj - ruj \\ 0 & 0 & 0 & pr(2h - e - j) \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$E_3 = \begin{bmatrix} 0 & t(e - a - 1) & tf + uh - au - bp - u & tg + ui + vj - av - bq - cr - v \\ 0 & 0 & p(h - e - 1) & pi + qj - eq - fr - q \\ 0 & 0 & 0 & r(j - h - 1) \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Of the ten brackets, three, $[E_2, E_4] = E_1$, $[E_3, E_4] = E_1$, $[E_4, E_5] = E_3 + E_4$, are satisfied by construction. Furthermore $[E_1, E_2] = 0$, $[E_1, E_3] = 0$, $[E_1, E_4] = 0$ identically. Of the remaining four brackets we shall need only the following:

$$[E_1, E_5] - 3E_1 = \begin{bmatrix} 0 & 0 & 0 & tpr(3e - a - 3h + j)(j - a - 3) \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$[E_3, E_5] - E_3 = \begin{bmatrix} 0 & t(e - a - 1)^2 & * & * \\ 0 & 0 & p(h - e - 1)^2 & * \\ 0 & 0 & 0 & r(j - h - 1)^2 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Considering E_1 , we see that $prt \neq 0$. Then looking at $[E_3, E_5] - E_3 = 0$ we see that we must have $e = a + 1$, $h = e + 1$, $j = h + 1$ and hence $3e - a - 3h + j = 0$ and hence $E_1 = 0$ which is a contradiction. ■

PROPOSITION 9.5. *Algebra $A_{5,38}$ has no representation in $\mathfrak{gl}(4, \mathbb{R})$*

Proof. Define $\{E_1, E_2, E_4, E_5\}$ as

$$E_1 = \begin{bmatrix} 0 & \alpha & \beta & \delta \\ 0 & 0 & \epsilon & \phi \\ 0 & 0 & 0 & \psi \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad E_2 = \begin{bmatrix} 0 & \rho & \sigma & \tau \\ 0 & 0 & \lambda & \mu \\ 0 & 0 & 0 & \theta \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$E_4 = \begin{bmatrix} a & b & c & d \\ 0 & e & f & g \\ 0 & 0 & h & i \\ 0 & 0 & 0 & j \end{bmatrix}, \quad E_5 = \begin{bmatrix} m & n & p & q \\ 0 & r & s & u \\ 0 & 0 & t & v \\ 0 & 0 & 0 & w \end{bmatrix}.$$

Then define $E_3 = [E_4, E_5]$ and we must have that $[E_1, E_4] = E_1$, $[E_2, E_5] = E_2$ and the remaining seven commutators are zero. Consulting the (3,4) entries of $[E_3, E_4]$ and $[E_3, E_5]$ we deduce that the (3,4)th entry of E_3 vanishes. Likewise the (1,2) and (2,3) entries of E_3 vanish.

Just as in $A_{5,22}$ there are seven normal forms for E_2 under change of basis. We work through each of these cases again suppressing the details of the very complicated matrices concerned. In the first case $\rho = 1$, $\lambda = 1$, $\theta = 1$, $\sigma = 0$, $\tau = 0$, $\mu = 0$ From $[E_2, E_5]$ we must have that $r = m + 1$, $s = n$, $u = p$, $t = r + 1$, $v = s$, $w = t + 1$ and from $[E_1, E_2]$ that $\epsilon = \alpha$, $\phi = \beta$, $\psi = \epsilon$. From $[E_1, E_5]$ we have that $\alpha = 0$, $\beta = 0$, $\delta = 0$ but now $E_1 = 0$.

In the second case (ii) $\rho = 1$, $\lambda = 1$, $\theta = 0$, $\sigma = 0$, $\tau = 0$, $\mu = 0$. Then $[E_2, E_5] = 0$ implies that $r = m + 1$, $s = n$, $u = 0$, $v = 0$, $t = r + 1$ and $[E_1, E_2] = 0$ that $\epsilon = \alpha$, $\phi = 0$, $\psi = 0$. Now $[E_1, E_5] = 0$ easily gives $\alpha = 0$, $\beta = 0$ and $[E_2, E_4] = 0$ that $e = a$, $f = b$, $g = 0$, $i = 0$, $h = e$. Finally the (1,3)th entry of $[E_3, E_5]$ implies that $c = 0$ and then E_1 and E_3 are proportional.

In the third case $\rho = 1$, $\lambda = 0$, $\theta = 1$, $\sigma = 0$, $\tau = 0$, $\mu = 0$. Then $[E_2, E_5] = 0$ implies that $r = m + 1$, $s = 0$, $u = p$, $w = t + 1$, from $[E_1, E_2] = 0$ that $\epsilon = 0$, $\phi = \beta$ and from $[E_1, E_5] = 0$ that $\alpha = 0$, $\psi = 0$. Next from $[E_2, E_4] = 0$ we find that $e = a$, $f = 0$, $g = c$, $j = h$ and from $[E_4, E_5] = E_3$ that $b = 0$, $i = 0$. Now the (1,4)th entry of $[E_2, E_3] = 0$ implies that $ap + ct - mc - ph = 0$. In order for E_3 not to vanish we must have that $cv + d - nc \neq 0$ and then from the the (1,4)th entries of $[E_3, E_4]$ and $[E_3, E_5]$ $h = a$, $t = m - 1$. Finally $[E_4, E_5] = E_3$ implies that $c = 0$, $d = 0$ and hence $E_3 = 0$.

In the fourth case $\rho = 0$, $\lambda = 0$, $\theta = 0$, $\sigma = 1$, $\tau = 0$, $\mu = 1$. From $[E_2, E_5] = 0$ we have that $t = m + 1$, $v = n$, $w = r + 1$, from $[E_1, E_2] = 0$ that $\psi = \alpha$ and from $[E_2, E_4] = 0$ that $h = a$, $i = b$, $j = e$. Now from $[E_3, E_5] = 0$ we find that $eu + nf + gm + g - rg - bs - ua = 0$ which can be used to simplify E_3 . The (1,3)th entry of $[E_3, E_5] = 0$ implies that $c = nf - bs$. Now $bu - nbs + n^2f + d - ng - pb \neq 0$ or else $E_3 = 0$ and then from the the (1,4) entries of $[E_3, E_4]$ and $[E_3, E_5]$ we deduce that $e = a$ and $r = m - 1$. Now from $[E_4, E_5] = E_3$ we have that $b = 0$, $d = 0$, $f = 0$, $g = 0$ which implies that $E_3 = 0$.

In the fifth case $\rho = 0$, $\lambda = 0$, $\theta = 0$, $\sigma = 1$, $\tau = 0$, $\mu = 0$. From $[E_2, E_5] = 0$ we have that $t = m + 1$, $v = 0$, from $[E_1, E_2] = 0$ that $\psi = 0$ and from $[E_2, E_4] = 0$ that $h = a$, $i = 0$. Now the (1, 3)th entries of $[E_3, E_5] = 0$ and $[E_1, E_5] = 0$ implies that $c = nf - bs$ and $\beta = n\epsilon - \alpha s$. Now we separate cases according as the (2, 4)th entry $eu + fv + gm + g - rg - si - ua$ of E_3 vanishes or not. In the first of these subcases the (1, 4)th entries of $[E_3, E_4] = 0$, $[E_3, E_5] = 0$, $[E_4, E_5] = E_3$ and $[E_1, E_4] = 0$ imply that $j = a$, $w = m$, $d = 0$ and $\delta = g\alpha - b\phi$. Now considering $[E_1, E_4] = 0$ we must have $e = a + 1$ or $e = a - 1$ or else $E_1 = 0$. However, now in either if these cases $[E_4, E_5] = E_3$ implies that $E_3 = 0$. Thus we may assume that $eu + fv + gm + g - rg - si - ua \neq 0$ and then the (1, 4)th entries of $[E_1, E_3] = 0$ and $[E_3, E_5] = 0$ imply that $\alpha = 0$ and $n = 0$. Then the (1, 2) and (2, 4) entries of $[E_3, E_4] = 0$ give $b = 0$ and $e = a$ and finally the (2, 4)th entry of $[E_3, E_5] = 0$ that $r = m$ and at this point we have that $[E_4, E_5] = E_3$ implies that $E_3 = 0$.

In the sixth case $\rho = 0$, $\lambda = 1$, $\theta = 0$, $\sigma = 0$, $\tau = 0$, $\mu = 0$. From $[E_2, E_5] = 0$ we have that $t = r + 1$, $n = 0$, $v = 0$, from $[E_1, E_2] = 0$ that $\alpha = 0$, $\psi = 0$ and from $[E_2, E_4] = 0$ that $b = 0$, $h = e$, $i = 0$. Now $[E_4, E_5] = E_3$ imply that $f = 0$ and the (2, 3)th entry of $[E_1, E_4] = E_1$ that $\epsilon = 0$. Now the (1, 3)th entries of $[E_3, E_4] = 0$ and $[E_3, E_5] = 0$ imply that the (1, 3)th entry $ap + cr + c - mc - pe = 0$ of E_3 vanishes. Suppose that $d \neq 0$. Then the (1, 4) entries of $[E_3, E_4] = 0$, $[E_3, E_5] = 0$ and $[E_4, E_5] = E_3$ imply that $j = a$, $w = m$, $d = 0$. Hence $d = 0$. Now comparing the (2, 4)th entries of $[E_3, E_4] = 0$ and E_3 we find that $j = e$ and from the (2, 4)th entry of $[E_3, E_5] = 0$ that $w = r$. Now, however, the (2, 4)th entry of $[E_4, E_5] = E_3$ implies that $E_3 = 0$.

In the seventh case $\rho = 0$, $\lambda = 0$, $\theta = 0$, $\sigma = 0$, $\tau = 1$, $\mu = 0$, consulting the (3, 4) entries of $[E_3, E_4]$ and $[E_3, E_5]$ we deduce that the (3, 4)th entry of $[E_3, E_4]$ vanish. Likewise the (1, 2) and (2, 3) entries of $[E_3, E_4]$ vanish. Now we cannot have that both the (1, 3) and (2, 4) entries of E_3 vanish or else E_2 and E_3 would be proportional. Appealing to Corollary 3.2 we may assume that the (1, 3)th entry of E_3 is not zero. Hence from the (1, 2) entries of $[E_3, E_4]$ and $[E_3, E_5]$ we deduce that $h = a$ and $t = m$. Now the (1, 2) entries of $[E_1, E_5]$ and $[E_4, E_5]$ imply that $\psi = 0$ and $i = 0$. Next the (1, 3) and (1, 4) entries of $[E_1, E_4]$ give that $\beta = f\alpha - b\epsilon$ and $\delta = g\alpha + fi\alpha - bi\epsilon - b\phi$ and the (1, 4)th entry of $[E_3, E_5]$ implies that $d = -(vbs - 2vnf + bu + cv - ngm - 2ng - pi - neu + nrg + nsi + nua)$. Consulting the (1, 2) and (2, 3) entries of $[E_1, E_4] = E_1$ we see that at least one of α and ϵ must be zero; hence from the (1, 2) and (2, 3) entries of $[E_1, E_5]$ we deduce that $r = m$. Next the

(2, 4) entries of $[E_3, E_5]$ and $[E_1, E_5]$ imply that $g = -(eu + fv - si - ua)$ and $\phi = -v\epsilon$.

Now we separate cases assuming first of all that $\alpha = 0$. Then $\epsilon \neq 0$ or else $E_1 = 0$. Now the (1, 3) entry of $[E_1, E_5]$ implies that $n = 0$ and the (2, 3) entry of $[E_1, E_4] - E_1$ and $[E_4, E_5] - E_3$ implies that $s = 0$. Now, however, $E_3 = 0$, a contradiction.

Finally suppose that $\epsilon = 0$ and $\alpha \neq 0$. Then the (1, 2) entry of $[E_1, E_4] - E_1$ gives that $e = a + 1$. The (1, 2) and (2, 3) entries of $[E_4, E_5] - E_3$ give $n = s = 0$. Now, however, again $E_3 = 0$. ■

Since $A_{5,39}$ is equivalent over \mathbb{C} to $A_{5,38}$ we deduce that

COROLLARY 9.6. $A_{5,39}$ has no representation in $\mathfrak{gl}(4, \mathbb{R})$.

REFERENCES

- [1] R. GHANAM, G. THOMPSON, Minimal matrix representations of four-dimensional Lie algebras, *Bull. Malays. Math. Sci. Soc. (2)* **36** (2) (2013), 343–349.
- [2] R. GHANAM, I. STRUGAR, G. THOMPSON, Matrix representations for low dimensional Lie algebras, *Extracta Math.* **20** (2) (2005), 151–184.
- [3] J. HUMPHREYS, “Introduction to Lie Algebras and Representation Theory”, Graduate Texts in Mathematics, vol. 9, Springer-Verlag, New York-Berlin, 1972.
- [4] N. JACOBSON, Schur’s Theorems on commutative matrices, *Bull. Amer. Math. Soc.* **50** (1944), 431–436.
- [5] Y. KANG, C. BAI, Refinement of Ado’s theorem in low dimensions and applications in affine geometry, *Comm. Algebra* **36** (1) (2008), 82–93.
- [6] G.N. MUBARAKZANOV, Classification of real Lie algebras in dimension five, *Izv. Vysshikh Uchebn. Zavedneii Mat.* **3** (34) (1963), 99–106.
- [7] J. PATERA, R.T. SHARP, P. WINTERNITZ, H. ZASSENHAUS, Invariants of real low dimension Lie algebras, *J. Mathematical Phys.* **17** (6) (1976), 986–994.
- [8] M. RAWASHDEH, G. THOMPSON, The inverse problem for six-dimensional codimension two nilradical Lie algebras, *J. Math. Phys.* **47** (11) (2006), 112901, 29 pp.